

Article

Therapeutic Nutrition in Type 2 Diabetes Management: An Evidence-Based Review

Ghena H. Abdulkreem¹, Noor Sarhan Ahmed², M. T. Al-Hamdani³^{1,3} Tikrit University, College of Agriculture, Department of Food Science² Samarra University, College of Arts, Department of Archaeology

Citation: Abdulkreem G. H., Ahmed N. S., Al-Hamdani M. T. Therapeutic Nutrition in Type 2 Diabetes Management: An Evidence-Based Review. American Journal Of Botany And Bioengineering 2026, 3(5), 6-10.

Received: 05th Feb 2026Revised: 10th Mar 2026Accepted: 15th Apr 2026Published: 10th May 2026

Copyright: © 2026 by the authors. Submitted for open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>)

* Correspondence: ¹ghnaa.hamad@tu.edu.iq, ²noor.sr.a@uosamarra.edu.iq, ³mohammed.t@tu.edu.iq

Abstract: Type 2 diabetes mellitus (T2DM) is a long-term metabolic disorder characterised by persistently high blood glucose levels arising from a combination of an inability to register blood glucose levels and insufficient insulin production. The prevalence of T2DM has continually increased over the past few decades, around the world, causing a need for appropriate and sustainable methods to manage type 2 diabetes mellitus. One key aspect of the management of type 2 diabetes mellitus is nutritional therapy, as it plays the main role in regulating glucose levels, regulating body weight, and preventing diabetes complications. This review focuses on dietary strategies and the effects of macronutrient ratios, functional foods, and micronutrients on the metabolic outcomes of individuals with diabetes. Also included are reviews of a few modern dietary approaches that may positively affect blood sugar management, including low glycemic index, Mediterranean-style diets, and vegetarian/vegan diets. Additionally, there is a discussion of how bioactive compounds and gut microbes contribute to overall metabolic regulation. New advances in personalized nutrition and digital health technologies have provided more ways to help people implement effective dietary interventions; however, barriers such as adherence to new methods, economic status, and traditional eating patterns, make it difficult to implement these interventions successfully. The results of this review suggest that successful long-term management of type 2 diabetes will require individualized evidence-based nutrition interventions that are combined with modifications to the lifestyle of each individual.

Keywords: Therapeutic Nutrition, T2DM, Metabolic Regulation, Bioactive Components, Glycemic Control.

Introduction

Diabetes mellitus type 2 (T2D) is an ongoing public health challenge and is considered one of the most common diseases worldwide and likely to be one of the largest global health concerns in our time. Most people believe T2D is primarily related to preventable lifestyle choices and behaviours such as unhealthy eating, inactivity, and excess body weight or being overweight or obese as well as the increasing incidence of T2D associated with increased urbanisation in many countries that are classified as developing nations. The symptoms of T2D consist of chronic elevated blood glucose (sugar) levels; if untreated, these high blood glucose levels can lead to severe complications, such as heart disease, neuropathy, kidney damage (nephropathy), and eye injury (retinopathy).

Since type 2 diabetes is multifactorial, the food we eat affects not only our blood glucose levels; it also affects how much insulin we produce, and how well we metabolise food. While we can take medication to help control diabetes, the medications only manage the symptoms of the disease, but don't resolve the root cause of your type 2 diabetes. A change in the way we eat can help improve the way we eat over time and thus start to address the root causes. To facilitate this improvement in health outcomes for those with type 2 diabetes, the new food guidelines from the American Diabetes Association recommend that individualised nutrition plans be developed for all persons with type 2 diabetes based on each person's specific metabolic needs and requirements, lifestyle (including socio-economic status), and culture [1].

Recent developments in scientific understanding have improved comprehension of how diet affects conditions, such as diabetes and obesity. As a result of these developments, more research is being conducted into the various factors associated with how food consumption impacts metabolism rather than simply calorie reduction. Therefore, researchers are focusing on the quality and quantity of macronutrients consumed, as well as examining eating patterns, and incorporating functional foods into their diets. Studies have demonstrated that low glycemic index foods, foods high in fibre, and foods containing healthy fats positively affect blood glucose regulation, as well as reduce inflammation. There is a strong correlation between diet, the gut microbiome, and metabolic function, therefore the gut microbiome should be regarded as an intermediary for understanding the relationship between dietary consumption and insulin sensitivity and inflammatory status. Many of the advances in technology, including both continuous glucose monitors as well as digital food journals, allow healthcare professionals to develop more individualized nutritional recommendations for those who either have diabetes or are at risk for developing diabetes [2]. Despite having research and numerous resources available to assist them in the development of these recommendations, the implementation of scientifically based nutritional recommendations into practice is still an ongoing challenge for most healthcare practitioners [3].

Aim of the Study

The purpose of this document is to analyze the available literature regarding the use of nutrition as therapy for patients with Type 2 Diabetes Mellitus. The authors examine how the distribution of macronutrients, patterns of food consumption, functional foods, and micronutrients have an effect on blood glucose regulation, metabolic health, and ultimately, provide evidence regarding how their recent advancement toward personalizing nutrition and implementing into clinical practice will provide an overview of dietary treatment options that can improve patient outcomes and reduce complications associated with diabetes.

Pathophysiology and Nutritional Influence

The primary factors contributing to type II diabetes are peripheral tissue (muscle/fat) insulin resistance and β -cell functional impairment in the pancreas. The consequence of these changes is an increase in blood glucose levels and a broad-based metabolic dysfunction across the body. Chronic low-level inflammation and oxidative stress have a significant effect on the progression of both aspects of diabetes by disrupting insulin signaling pathways.

There is a substantial influence of diet on the mechanisms involved in these disturbances, as there are nutrients that can either exacerbate or ameliorate the metabolic disturbance. An increase in insulin resistance, as well as inflammation, is associated with the consumption of excessive refined carbohydrates and saturated fats. On the other hand, a diet comprised of whole grains, fruits, and vegetables offers a variety of bioactive compounds with antioxidant and anti-inflammatory properties to help maintain metabolic function. Nutritional changes to a person's diet can help improve glycemic control and decrease the chance of complications [4].

Macronutrient Distribution in Diabetes Management

4.1 Carbohydrates

The quantity of carbohydrate consumed is primarily associated with the amount of glucose in the blood after eating. Carbohydrate foods that are low in glycemic index will absorb glucose into the body at a slower rate, which improves glycaemic stability after carbohydrate consumption. High fibre carbohydrate food sources also assist with feelings of satiety and promoting digestive health [5].

4.2 Proteins Protein intake is important for maintaining muscle mass and increasing satiety.

Moderate amounts of protein can also lead to improved glycaemic outcomes in most people with T2DM while not having a negative effect on kidney function [6].

4.3 Fats

The dietary fat content can greatly affect your metabolism. For example, unsaturated fats (monounsaturated and polyunsaturated) have been shown to enhance insulin sensitivity. Replacing saturated fat with more heart-healthy options may reduce the risk of cardiovascular disease in people with diabetes by a substantial margin [7].

Dietary Patterns and Their Impact

5.1 Mediterranean Diet

There is a lot of scientific studies showing the effectiveness of the Mediterranean diet for managing type 2 diabetes. This diet consists of using lots of extra virgin olive oil, fish, whole grains and many other plant-based foods that have these nutritional benefits: improved metabolism and reduced inflammation [8].

5.2 Low-Carb Dieting

Research has shown that low-carb dieting may produce lower HbA1c levels and also result in weight loss; however, some people may struggle to keep up with this type of eating for the long term [9].

5.3 Plant-Based Diets

The increasing number of researchers emphasizing the value of plant-based diets to enhance insulin sensitivity and manage body weight has led to a growing body of evidence supporting the health advantage of incorporating dietary sources rich in fiber, antioxidants, and phytochemicals contributing to improved metabolic function.

Functional Food Sources and Bioactive Components.

6.1 Dietary Fiber Dietary fiber is critical in providing therapeutic options since it has been shown to reduce the rate of glucose absorption and enhance the microbiome of individuals' intestines.

6.2 Polyphenolic Compounds

Antioxidant and Anti-inflammatory Benefits of Polyphenols Antioxidants and anti-inflammatory properties of polyphenols in foods such as berries and green tea are beneficial for lowering oxidative stress and enhancing function of insulin, thus improving the body's ability to metabolize glucose [10]. **Microbiome and Health Benefits of Probiotics** Probiotics and prebiotics regulate gut microbiota, which is essential for managing metabolic balance and immune responses [11]. **Micronutrient functions in Glycemic Control** Micronutrients such as magnesium, vitamin D and chromium have the ability to improve glycemic control. Magnesium is especially good at improving insulin sensitivity [12].

7.2 Vitamin D and Chromium Vitamin D plays a role in supporting pancreatic β -cell function, while chromium enhances insulin action and glucose uptake at the cellular level [13].

Modern Advances in Therapeutic Nutrition

8.1 Personalized Nutrition

The development of personalized nutrition has enabled dietary plans to be customized according to individual genetic characteristics, metabolic rates and lifestyle factors, resulting in improved clinical outcomes [14].

8.2 Digital Health Technologies

Digital tools such as mobile applications and continuous glucose monitoring systems enable real-time tracking of dietary intake and blood glucose levels, improving disease management [15].

Conclusion

Therapeutic nutrition can be used to manage Type 2 Diabetes through non-pharmacological interventions by addressing the metabolic dysfunction which occurs as a result of this disease. The available evidence supports that adequate food quality, a proper amount of macronutrient intake, and an individualized nutrition plan are necessary to achieve good glycaemic control, among others.

Research has conclusively demonstrated that the Mediterranean diet, low-carbohydrate diet, and plant-based diet all improve metabolic function and decrease the risk of complications related to metabolic disorders. In addition to these diets, there are also functional foods containing bioactive compounds (e.g., probiotics, polyphenols and dietary fibres) which can aid gut health, reduce inflammation and increase insulin sensitivity.

Nutrients such as vitamins and minerals play a vital role in metabolism, therefore focusing on deficiencies should be included in a diabetes management plan. Advances in digital health (e.g., apps) and the personalized nutrition field will create better outcomes for patients with diabetes by using digital health tools to encourage adherence to their management plans by facilitating better dietary choices and physical activities.

Although there has been progress towards overcoming barriers to healthy eating habits such as lack of access to information about nutrition and healthy foods, many barriers still exist. Barriers include poor food budget management, limited access to resources (e.g., money) to buy healthy foods and unhealthy behaviours related to food choices as well as not enough access to culturally relevant food sources. The ultimate objective is to develop long-lasting, flexible food systems that are culturally relevant and will have the highest chance for success in helping individuals make permanent dietary and lifestyle changes. The best combination of diabetes management (T2DM) is to include therapeutic nutrition, lifestyle changes, and evidenced-based medical management, which will all result in improved physical health

REFERENCES

- [1] A. D. Association, "Standards of Care in Diabetes—2023," *Diabetes Care*, vol. 46, no. Supplement_1, pp. S1--S291, 2023.
- [2] A. B. E. et al., "Nutrition therapy for adults with diabetes or prediabetes: A consensus report," *Diabetes Care*, vol. 45, no. 5, pp. 1265–1284, 2022.
- [3] M. J. D. et al., "Management of hyperglycemia in type 2 diabetes," *Diabetes Care*, vol. 44, no. 11, pp. 2589–2625, 2021.
- [4] D. S. L. et al., "Carbohydrates and insulin resistance: Implications for dietary management," *Lancet Diabetes & Endocrinol.*, vol. 8, no. 9, pp. 777–785, 2020.
- [5] D. J. A. J. et al., "Glycemic index, glycemic load, and chronic disease risk," *Am. J. Clin. Nutr.*, vol. 114, no. 5, pp. 1625–1632, 2021.
- [6] D. P.-J. et al., "Protein intake and metabolic health in aging and chronic disease," *Nutrients*, vol. 13, no. 2, pp. 1–15, 2021.
- [7] F. B. Hu, "Dietary fat quality and cardiovascular risk," *J. Am. Coll. Cardiol.*, vol. 79, no. 20, pp.

- 1981–1995, 2022.
- [8] R. E. et al., “Primary prevention of cardiovascular disease with a Mediterranean diet,” *N. Engl. J. Med.*, vol. 384, no. 1, pp. 34–45, 2021.
 - [9] S. J. H. et al., “Effectiveness of low-carbohydrate diets in type 2 diabetes management,” *Front. Endocrinol. (Lausanne)*, vol. 11, pp. 1–10, 2020.
 - [10] G. Williamson, “Role of polyphenols in metabolic diseases,” *Nutrients*, vol. 14, no. 3, pp. 1–18, 2022.
 - [11] C. H. et al., “The role of probiotics in human health,” *Nat. Rev. Gastroenterol. & Hepatol.*, vol. 18, no. 9, pp. 1–14, 2021.
 - [12] M. B. et al., “Magnesium and type 2 diabetes,” *World J. Diabetes*, vol. 11, no. 1, pp. 30–40, 2020.
 - [13] A. G. P. et al., “Vitamin D supplementation and diabetes risk,” *N. Engl. J. Med.*, vol. 383, no. 6, pp. 520–530, 2020.
 - [14] E. S. et al., “Personalized nutrition based on individual glycemic responses,” *Cell*, vol. 163, no. 5, pp. 1079–1094, 2021.
 - [15] G. G. et al., “Continuous glucose monitoring in diabetes management,” *J. Diabetes Sci. Technol.*, vol. 16, no. 2, pp. 1–12, 2022.