

The Role of the Immune System in Maintaining the Constancy of the Internal Environment of the Body

Ismatova Marguba Shaukatovna

Senior Teacher of the Department of Physiology, Samarkand State Medical University

Nasimova Diyora Ikromovna

Student of Faculty of Medicine No. 1, Group 216, Samarkand State Medical University

Received: 2024, 26, Aug

Accepted: 2025, 28, Sep

Published: 2025, 29, Oct

Copyright © 2025 by author(s) and BioScience Academic Publishing. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).



Open Access

<http://creativecommons.org/licenses/by/4.0/>

Annotation: The immune system plays a fundamental role in preserving the stability and constancy of the body's internal environment, a process known as homeostasis. It protects the organism from harmful pathogens, foreign substances, and internal abnormalities such as malignant cells. Through a complex network of cells, tissues, and signaling molecules, the immune system continuously monitors the body, identifying and eliminating threats while maintaining tolerance to self-antigens. This article explores the physiological mechanisms of immune regulation in maintaining homeostasis, focusing on innate and adaptive immunity, cytokine signaling, and immune tolerance. The research highlights how disturbances in immune balance can lead to pathological conditions such as autoimmune diseases, allergies, and immunodeficiency, emphasizing the immune system's central role in overall health and homeostatic equilibrium.

The immune system is the key physiological mechanism that ensures the balance and stability of the human body by protecting it from external and internal threats. It performs a complex network of functions that include recognition, neutralization, and elimination of pathogenic microorganisms, abnormal cells, and toxic

substances while maintaining tolerance toward the body's own tissues. This article discusses how the immune system maintains homeostatic balance, analyzes the functional interaction between innate and adaptive immunity, and describes how the dysregulation of immune responses leads to various pathological processes. The immune system, acting as a self-regulating defense mechanism, constantly monitors the internal environment, adjusts immune responses according to the body's needs, and restores equilibrium after infection, injury, or stress.

Keywords: Immune system, homeostasis, internal environment, immune response, cytokines, innate immunity, adaptive immunity, immune tolerance, inflammation, autoimmunity.

Introduction:

The human body is constantly exposed to various environmental factors, microorganisms, and internal metabolic changes that can disrupt its stability. The concept of homeostasis, first proposed by Claude Bernard and later expanded by Walter Cannon, describes the ability of living organisms to maintain a stable internal environment despite external fluctuations. The immune system is one of the primary regulatory systems ensuring this constancy. Its main function is to detect and neutralize potentially harmful agents while preserving the integrity of normal body tissues. The immune system operates as a highly coordinated network involving the bone marrow, thymus, spleen, lymph nodes, and peripheral immune cells such as macrophages, lymphocytes, and dendritic cells. These components act together to recognize antigens, trigger appropriate immune responses, and restore balance after infection or injury. The maintenance of immune homeostasis requires a delicate balance between activation and suppression. Overactivation can result in autoimmune reactions, whereas insufficient activation can lead to increased susceptibility to infections and tumors. Thus, understanding the role of the immune system in maintaining internal constancy is essential for explaining the pathophysiology of many diseases and developing new therapeutic strategies.

The human organism is an open biological system that constantly interacts with the external environment. For survival, it must maintain a stable internal state despite external changes. This physiological stability, known as homeostasis, is achieved through the coordinated activity of several systems—nervous, endocrine, and immune. Among them, the immune system plays a central role because it defends the body against pathogenic microorganisms, repairs damaged tissues, and regulates inflammatory reactions. It acts as a biological “security system,” identifying all that is foreign and harmful, while protecting self-cells and maintaining structural and functional harmony. The immune system consists of organs and cellular elements such as the thymus, spleen, lymph nodes, bone marrow, macrophages, lymphocytes, and various signaling molecules. Together, these components work to detect threats and initiate immune responses. However, the immune system's role extends beyond defense—it also contributes to tissue homeostasis, wound healing, and elimination of dead or mutated cells. Any disruption in immune regulation can cause severe consequences: hyperactivation may lead to autoimmunity,

while immune suppression can result in infections and malignancies. Thus, the immune system is both a protector and a regulator, ensuring that all physiological processes remain in dynamic balance.

Materials and Methods:

This study was based on an analytical review of scientific literature and clinical observations conducted at Samarkand State Medical University between 2022 and 2024. The research included evaluation of immune parameters in healthy individuals and patients with immune-related disorders. Laboratory analyses focused on the quantification of white blood cells, lymphocyte subpopulations (CD4+, CD8+, B cells, NK cells), and cytokine levels (IL-2, IL-6, TNF- α , and interferon- γ). Blood samples were collected and processed using flow cytometry and enzyme-linked immunosorbent assay (ELISA). Comparative assessment of immune responses under normal and pathological conditions was carried out to determine how immune activity contributes to maintaining homeostasis. Additionally, data on inflammatory markers such as C-reactive protein (CRP) and erythrocyte sedimentation rate (ESR) were analyzed to evaluate the systemic response to immune disturbances.

Results:

The analysis revealed that in healthy individuals, immune cell populations and cytokine levels remain within narrow physiological ranges, reflecting stable immune homeostasis. Any deviation from these parameters correlated with either hyperactivation or suppression of immune function. For example, elevated levels of proinflammatory cytokines such as IL-6 and TNF- α were associated with systemic inflammatory responses, leading to disturbances in temperature regulation, metabolism, and vascular tone. Conversely, low cytokine activity was linked to increased vulnerability to infections and slower wound healing. The study also confirmed that adaptive immune components, particularly T regulatory (Treg) cells, play a vital role in maintaining self-tolerance and preventing autoimmune processes. In patients with immune dysregulation, reduced Treg activity was observed, resulting in the breakdown of tolerance and development of autoimmune pathologies such as rheumatoid arthritis and systemic lupus erythematosus. Overall, the immune system's balance between proinflammatory and anti-inflammatory mechanisms proved crucial for preserving the body's internal equilibrium.

The functional activity of the immune system directly determines the stability of the internal environment. Experimental and clinical observations show that immune responses are finely tuned to maintain equilibrium. When a pathogen enters the body, innate immunity acts first, producing inflammation to isolate and destroy the invader. This early response activates adaptive immunity, which forms specific antibodies and immune memory cells to prevent future infections. The balance between these two components—innate and adaptive—is essential for sustained homeostasis. In a well-functioning organism, proinflammatory and anti-inflammatory cytokines are in constant equilibrium, ensuring controlled immune activation and timely resolution of inflammation. The immune system not only fights infections but also monitors internal cellular health through immune surveillance. Natural killer (NK) cells and cytotoxic T lymphocytes detect and eliminate transformed or cancerous cells, preventing their proliferation. In addition, the immune system supports the regeneration of tissues by releasing growth factors that promote healing and remodeling. Disturbances in this system lead to specific outcomes: excessive immune activity causes chronic inflammation and tissue damage, whereas reduced immunity increases susceptibility to disease. The data confirm that immune stability is the foundation of all physiological harmony in the human body.

Discussion:

The findings underscore that the immune system is not only a defense mechanism but also a dynamic regulator of homeostasis. Innate immunity acts as the first barrier, providing rapid, nonspecific defense through phagocytosis, complement activation, and inflammation. This initial

response is followed by adaptive immunity, which ensures antigen-specific memory and long-term protection. Cytokines, chemokines, and growth factors function as signaling molecules that coordinate interactions between immune cells and other physiological systems such as the endocrine and nervous systems. Immune surveillance constantly monitors the body for the presence of pathogens and abnormal cells, removing them before they can disrupt internal stability. Importantly, immune tolerance mechanisms—both central and peripheral—prevent self-reactivity, thereby avoiding unnecessary tissue damage. Disturbances in immune regulation can result in pathological conditions that compromise homeostasis. Chronic inflammation, for instance, leads to metabolic imbalances and tissue destruction, while immunodeficiency reduces the body's ability to adapt to stress and infection. Moreover, immune-endocrine interactions demonstrate that hormonal fluctuations can influence immune function, indicating the integrative nature of homeostatic regulation. Maintaining a balanced immune response is therefore essential for overall physiological stability and longevity.

The maintenance of homeostasis depends on the precise regulation of immune responses. The immune system communicates closely with the endocrine and nervous systems, forming an integrated network of regulation known as the neuro-immune-endocrine axis. Hormones such as cortisol, adrenaline, and cytokines mediate feedback mechanisms that either stimulate or suppress immune activity depending on the situation. During infection or injury, the immune system increases its activity to neutralize harmful agents; once the threat is removed, anti-inflammatory signals restore the internal balance. This adaptability allows the body to respond effectively without damaging its own tissues. However, modern lifestyles—chronic stress, poor diet, pollution, and lack of physical activity—can disturb immune regulation, leading to systemic inflammation and metabolic disorders. The immune system's ability to distinguish "self" from "non-self" is another crucial factor in maintaining constancy. This recognition prevents the destruction of body cells and ensures tolerance to self-antigens. Autoimmune diseases develop when this recognition fails, illustrating how central immune control is to physiological stability. Research also demonstrates that the gut microbiota plays an important role in regulating immune function. A balanced intestinal flora strengthens immune tolerance and supports metabolic health, while dysbiosis triggers inflammation and immune imbalance. Hence, the immune system serves not only as a defense mechanism but also as an adaptive regulator that keeps the body in a state of dynamic equilibrium.

Conclusion:

The immune system plays a central and multifaceted role in maintaining the constancy of the body's internal environment. Through continuous surveillance, recognition, and elimination of harmful agents, it ensures protection while preserving tissue integrity. Proper coordination between innate and adaptive immunity maintains equilibrium and allows the body to adapt to changing external and internal conditions. Any imbalance in immune regulation can disrupt homeostasis, leading to diseases ranging from infections to autoimmune and inflammatory disorders. Strengthening immune function through proper nutrition, physical activity, vaccination, and stress management contributes to maintaining homeostatic balance and overall health. Understanding the physiological and molecular basis of immune homeostasis provides the foundation for new preventive and therapeutic approaches in modern medicine.

The immune system is indispensable in maintaining the constancy of the internal environment and ensuring the organism's survival. By recognizing and eliminating foreign agents, regulating inflammatory processes, and promoting tissue repair, it continuously preserves the body's functional integrity. The harmony between immune activation and suppression is vital for sustaining homeostasis; any deviation from this balance can result in disease. The immune system interacts with other physiological systems through a network of signaling molecules that maintain coordination and adaptability. Strengthening immune resilience through a balanced lifestyle, proper nutrition, sufficient rest, and vaccination supports overall health and long-term stability. In essence, the immune system is not merely a defense barrier but a central regulatory

force that sustains the equilibrium of all vital processes within the human body, ensuring its ability to adapt and thrive under ever-changing conditions.

References:

1. Abbas, A. K., Lichtman, A. H., & Pillai, S. (2022). *Cellular and Molecular Immunology*. Elsevier.
2. Murphy, K., & Weaver, C. (2017). *Janeway's Immunobiology*. Garland Science.
3. Alberts, B. et al. (2022). *Molecular Biology of the Cell*. Garland Science.
4. Medzhitov, R., & Janeway, C. A. (2000). Innate immune recognition. *Annual Review of Immunology*, 18, 217–242.
5. Chaplin, D. D. (2010). Overview of the immune response. *Journal of Allergy and Clinical Immunology*, 125(2), S3–S23.
6. Nicholson, L. B. (2016). The immune system. *Essays in Biochemistry*, 60(3), 275–301.
7. Iwasaki, A., & Medzhitov, R. (2015). Control of adaptive immunity by the innate immune system. *Nature Immunology*, 16(4), 343–353.
8. Kotas, M. E., & Medzhitov, R. (2015). Homeostasis, inflammation, and disease susceptibility. *Cell*, 160(5), 816–827.
9. O'Shea, J. J., & Murray, P. J. (2008). Cytokine signaling modules in inflammatory responses. *Immunity*, 28(4), 477–487.
10. Hotamisligil, G. S. (2017). Inflammation, metaflammation, and immunometabolic disorders. *Nature*, 542(7640), 177–185.